


Edited by
Makoto M. Watanabe, Ken-ichiro Suzuki & Tatsuji Seki

Makoto M. Watanabe, Ken-ichiro Suzuki & Tatsuji Seki



Innovative Roles of Biological Resource Centers

Proceedings of the Tenth International Congress for Culture Collections
Tsukuba, Japan, 10-15 October 2004

Edited by
Makoto M. Watanabe
Ken-ichiro Suzuki
Tatsuji Seki

Japan Society for Culture Collections &
World Federation for Culture Collections

Biosynthesis, detection and toxicology of ochratoxins - optimisation of production

Luis Abrunhosa^{1*}, Lucia Santos², Armando Venancio¹

¹Centro de Engenharia Biologica, Universidade do Minho, Campus de Gualtar, 4710-057 Braga, Portugal

²LEPA, Departamento de Engenharia Quimica, Faculdade de Engenharia da Universidade do Porto, Rua Dr. Roberto Frias 4200-465 Porto, Portugal

Author for correspondence: email <luisjap@deb.uminho.pt>

Ochratoxin A (OTA) is a very well known mycotoxin found in several food commodities for which maximum limits are being discussed in EC in order to produce appropriate regulations. OTA is one of several ochratoxins produced by *Aspergillus* and *Penicillium* species. All the compounds in this group have a molecular structure very similar to OTA and some were already isolated from natural substrates. Several of these compounds such as ochratoxin, methyl and ethyl ester of ochratoxin A, 4-*R* and *S*-hydroxyochratoxin A, 10-hydroxyochratoxin A and ochratoxin A open lactone are commercially unavailable. However, they can be easily synthesized through OTA modification. With the main objective of its application on further research works, OTA production, isolation and purification has been optimised from an *A. alliaceus* strain grown on wheat medium. Synthesis and purification of some OTA derivatives has been achieved and an HPLC method for their detection was optimised. Data about their production by several species of *Aspergillus* will be presented. The toxicological properties of

Mycology

ochratoxins are still not very clear and a future EC safety limit for OTA will depend on e.g., a better clarification of its carcinogenicity. Could OTA derivatives play a role here?